

Marine dinoflagellate cysts as tools in palaeoecology

Mertens Kenneth Neil

Research Unit Palaeontology, Ghent University, Krijgslaan 281, S8, B-9000 Gent, Belgium
E-mail: Kenneth.mertens@ugent.be

Dinoflagellates form a large group of flagellate protists with about 1,700 marine species (free-living as well as benthic) and about 220 described from freshwater (Taylor *et al.*, 2008). Some dinoflagellates produce a resting stage, called the cyst or dinocyst, as a part of their life cycle (Dale, 1983). These cysts often are well-preserved as microfossils in the sediments, depending on their geochemical composition (e.g. Bogus *et al.*, 2014). Dinocysts are excellent palaeoecological indicators: both dinocyst assemblages and their morphological variations have been shown to reflect changes over time in temperature, salinity, ice cover and productivity as well as industrial pollutants and coastal proximity (e.g. Dale, 1996).

Here I will show several examples of the palaeoecological importance of these dinocysts. Firstly I will show how the morphological variations can be used to reconstruct salinity in the Holocene Black Sea (Mertens *et al.*, 2012). Secondly, I will highlight the spectacular discovery of living cysts of *Dapsilidinium pastielsii* in the Indo-Pacific Warm Pool, which served as a refuge during glacial times (Mertens *et al.* 2014). Thirdly, I will briefly discuss the importance of cysts in the toxic species *Pyrodinium bahamense* (Mertens *et al.*, 2015).

References

- Bogus K., K.N. Mertens, J. Lauwaert, I.C. Harding, H. Vrielinck, K.A.F. Zonneveld, and G.J.M. Versteegh. 2014. Differences in the chemical composition of organic-walled dinoflagellate resting cysts from phototrophic and heterotrophic dinoflagellates. *Journal of Phycology* 50:254–266.
- Dale B. 1983. Dinoflagellate resting cysts: “benthic plankton”. *In*: Survival strategies of the algae. Fryxell G.A.(Eds). Cambridge University Press, London and New York.
- Dale B. 1996. Dinoflagellate cyst ecology: modelling and geological applications. p.1249–1275. *In*: Palynology: principles and applications. Vol. 3. Jansonius J., D.C. McGregor (Eds). AASP Foundation, Dallas, TX.
- Mertens K.N., L.R. Bradley, Y. Takano, P.J. Mudie, F. Marret, A.E. Aksu, R.N. Hiscott, T.J. Verleye, E.A. Mousing, L.L. Smyrnova, S. Bagheri, M. Mansor, V. Pospelova, K. Matsuoka. 2012. Quantitative estimation of Holocene surface salinity variation in the Black Sea using dinoflagellate cyst process length. *Quaternary Science Reviews* 39:45–59.
- Mertens K.N., Y. Takano, M.J. Head, and K. Matsuoka. 2014. Living fossils in the Indo-Pacific warm pool: a refuge for thermophilic dinoflagellates during glaciations. *Geology* 42(6):531–534.
- Mertens K.N., J. Wolny, C. Carbonell-Moore, K. Bogus, M. Ellegaard, A. Limoges, A. de Vernal, P. Gurdebeke, T. Omura, A. Mohd, A. Al-Muftah, and K. Matsuoka. 2015. Taxonomic re-examination of the toxic armoured dinoflagellate *Pyrodinium bahamense* Plate 1906: can morphology or LSU sequencing separate *P. bahamense* var. *compressum* from var. *bahamense*? *Harmful Algae* 41:1–24.
- Taylor F.J.R., M. Hoppenrath, and J.F. Saldarriaga. 2008. Dinoflagellate diversity and distribution. *Biodiversity and Conservation* 17:407–418.